

RESPONSE

Response to concerns raised about the likelihood of protected areas serving as steppingstones for species responding to climate change

Sean A. Parks¹  | Lisa M. Holsinger¹ | John T. Abatzoglou²  | Caitlin E. Littlefield³  | Katherine A. Zeller¹

¹Aldo Leopold Wilderness Research Institute, Rocky Mountain Research Station, US Forest Service, Missoula, Montana, USA

²Management of Complex Systems, University of California, Merced, California, USA

³Conservation Science Partners, Truckee, California, USA

Correspondence

Sean A. Parks, Aldo Leopold Wilderness Research Institute, Rocky Mountain Research Station, US Forest Service, 790 E. Beckwith Ave., Missoula, MT 59801, USA.

Email: sean.parks@usda.gov

Keywords: climate change, climate change vulnerability, climate connectivity, climate connectivity failure, climate-wise connectivity, protected areas, species range shifts

Our recent contribution to Global Change Biology found that, globally, two-thirds of protected area units and over half of protected land area are unlikely to serve as steppingstones for many species under a warming climate (Parks et al., 2023). In response, Gillingham and Thomas (2023) point to previous evidence that protected areas have acted as steppingstones based on three studies from Great Britain that examined range shifts.

The first study found that 98% of the colonizing species analyzed ($n=251$) were more reliant on protected areas than unprotected land (Thomas et al., 2012). The second study found that 62% of the butterfly and odonate (dragonflies and damselflies) species analyzed ($n=99$) were more abundant in protected areas in newly colonized regions compared to the unprotected matrix (Gillingham et al., 2015). The third study found that bird species colonizing Great Britain usually established first in protected areas before expanding to unprotected lands (Hiley et al., 2014). Although these studies underscore the importance of PAs in helping species adapt to climate change—an importance we fully acknowledge—our research (Parks et al., 2023) provides nuance into factors that may impede successful range shifts among protected areas.

It is worth noting that a large proportion of the species in Thomas et al. (2012) and all species in Gillingham et al. (2015) and

Hiley et al. (2014) have wings and can therefore fly, which means many of these species are highly mobile. As pointed out by Parks et al. (2023), such species are not likely limited by dispersal and, by inference, protected areas are more likely to serve as steppingstones for such species. Other species and lifeforms (such as plants) often have more restrictive dispersal abilities, so although “most species” in Gillingham and Thomas (2023) may have used protected areas for range shifts, this by no means equates to most species overall. Moreover, our study looks ahead to a period when temperatures are likely to rise at a greater rate than experienced by Great Britain during the period examined by Gillingham and Thomas (2023). Even in the absence of protected areas that harbor higher levels of biodiversity compared to the unprotected matrix (Brodie et al., 2023), others have concluded that climate change velocity exceeds the dispersal abilities of many species (Burrows et al., 2014) and human land use hinders the ability of many species to track climate change (Robillard et al., 2015).

Parks et al. (2023) clearly recognized that the biodiversity at a given site comprises innumerable taxa, with each species having variable dispersal abilities, sensitivities to human land uses, and climatic tolerances. As we did not claim that *all* protected areas were unlikely to serve as steppingstones for *all* species, we stand by our

Lisa M. Holsinger, Retired.

This article is a Response to the Letter by Gillingham & Thomas, <https://doi.org/10.1111/gcb.16941>, which was related to the paper of Parks et al, <https://doi.org/10.1111/gcb.16629>

Published 2023. This article is a U.S. Government work and is in the public domain in the USA.

overall finding that protected areas are unlikely to serve as steppingstones for a large number of species under a warming climate. We thank Gillingham and Thomas (2023) for an interesting debate.

ACKNOWLEDGEMENTS

This research was supported in part by the USDA Forest Service. The findings and conclusions in this publication are those of the authors and should not be construed to represent any official USDA or US Government determination or policy.

ORCID

Sean A. Parks  <https://orcid.org/0000-0002-2982-5255>

John T. Abatzoglou  <https://orcid.org/0000-0001-7599-9750>

Caitlin E. Littlefield  <https://orcid.org/0000-0003-3771-7956>

REFERENCES

- Brodie, J. F., Mohd-Azlan, J., Chen, C., Wearn, O. R., Deith, M. C. M., Ball, J. G. C., Slade, E. M., Burslem, D. F. R. P., Teoh, S. W., Williams, P. J., Nguyen, A., Moore, J. H., Goetz, S. J., Burns, P., Jantz, P., Hakkenberg, C. R., Kaszta, Z. M., Cushman, S., Coomes, D., ... Luskin, M. S. (2023). Landscape-scale benefits of protected areas for tropical biodiversity. *Nature*, 620(7975), 807–812. <https://doi.org/10.1038/s41586-023-06410-z>
- Burrows, M. T., Schoeman, D. S., Richardson, A. J., Molinos, J. G., Hoffmann, A., Buckley, L. B., Moore, P. J., Brown, C. J., Bruno, J. F., Duarte, C. M., Halpern, B. S., Hoegh-Guldberg, O., Kappel, C. V., Kiessling, W., O'Connor, M. I., Pandolfi, J. M., Parmesan, C., Sydeman, W. J., Ferrier, S., ... Poloczanska, E. S. (2014). Geographical limits to species-range shifts are suggested by climate velocity. *Nature*, 507(7493), 492–495. <https://doi.org/10.1038/nature12976>
- Gillingham, P., & Thomas, C. (2023). Protected areas do already act as steppingstones for species responding to climate change. *Global Change Biology*, <https://doi.org/10.1111/gcb.16941>
- Gillingham, P. K., Alison, J., Roy, D. B., Fox, R., & Thomas, C. D. (2015). High abundances of species in protected areas in parts of their geographic distributions colonized during a recent period of climatic change. *Conservation Letters*, 8(2), 97–106. <https://doi.org/10.1111/conl.12118>
- Hiley, J. R., Bradbury, R. B., & Thomas, C. D. (2014). Introduced and natural colonists show contrasting patterns of protected area association in UK wetlands. *Diversity and Distributions*, 20(8), 943–951. <https://doi.org/10.1111/ddi.12219>
- Parks, S. A., Holsinger, L. M., Abatzoglou, J. T., Littlefield, C. E., & Zeller, K. A. (2023). Protected areas not likely to serve as steppingstones for species undergoing climate-induced range shifts. *Global Change Biology*, 29(10), 2681–2696. <https://doi.org/10.1111/gcb.16629>
- Robillard, C. M., Coristine, L. E., Soares, R. N., & Kerr, J. T. (2015). Facilitating climate-change-induced range shifts across continental land-use barriers. *Conservation Biology*, 29(6), 1586–1595. <https://doi.org/10.1111/cobi.12556>
- Thomas, C. D., Gillingham, P. K., Bradbury, R. B., Roy, D. B., Anderson, B. J., Baxter, J. M., Bourn, N. A. D., Crick, H. Q. P., Findon, R. A., Fox, R., Hodgson, J. A., Holt, A. R., Morecroft, M. D., O'Hanlon, N. J., Oliver, T. H., Pearce-Higgins, J. W., Procter, D. A., Thomas, J. A., Walker, K. J., ... Hill, J. K. (2012). Protected areas facilitate species' range expansions. *Proceedings of the National Academy of Sciences*, 109(35), 14063–14068. <https://doi.org/10.1073/pnas.1210251109>

How to cite this article: Parks, S. A., Holsinger, L. M., Abatzoglou, J. T., Littlefield, C. E., & Zeller, K. A. (2023). Response to concerns raised about the likelihood of protected areas serving as steppingstones for species responding to climate change. *Global Change Biology*, 00, 1–2. <https://doi.org/10.1111/gcb.16940>